

Path integral representation of the evolution operator for the Dirac equation

Alexander S. Lukyanenko*

*Department of Experimental Physics, St. Petersburg State Polytechnical University,
Polytekhnicheskaya 29, 195251, St. Petersburg, Russia*

Inna A. Lukyanenko†

*Department of Mathematics, St. Petersburg State University,
Universitetski prospect 28, Stary Petergof, 198504, St. Petersburg, Russia*

A path integral representation of the evolution operator for the four-dimensional Dirac equation is proposed. A quadratic form of the canonical momenta regularizes the original representation of the path integral in the electron phase space. This regularization allows to obtain a representation of the path integral over trajectories in the configuration space, i.e. in the Minkowsky space. This form of the path integral is useful for the formulation of perturbation theory in an external electromagnetic field.

PACS numbers:

Keywords:

The formulation of quantum mechanics in terms of complex amplitudes and its concentrated representation in terms of the Feynman path integral in a configuration space became very useful for the further development of quantum theory. In particular, this approach is the most suitable for the formulation of perturbation theory rules. Moreover, it is difficult to formulate the quantum theory of gauge fields without the Feynman path integral. However, one of the oldest parts of quantum theory, the relativistic quantum mechanics of an electron, which is based on the Dirac equation, has no path integral representation. Feynman has attempted to solve the problem¹. He hoped to understand the nature of the spin by use of this representation. He obtained a solution for an electron in one dimension (1D). In this case two components of an electron wave function are related to two possible ways to reach the end point of a path. For an electron in three dimensions (3D) such a clear solution was not obtained. In this paper we return to the problem of a path integral representation of solutions of the Dirac equation. We consider this representation as a tool for the formulation of perturbation theory rules for an evolution operator of the Dirac equation in an external electromagnetic field.

The Dirac equation for an electron in an external electromagnetic field described by the four-vector potential $A_\mu = (\varphi, A_i)$ has a form²:

$$i\hbar \frac{\partial \psi}{\partial t} = \hat{h}\psi \equiv c\hat{\alpha}_k \left(\frac{\hbar}{i} \nabla_k - \frac{e}{c} A_k \right) \psi + \hat{\beta} mc^2 \psi + e\varphi \psi \quad (1)$$

Here ψ is the Dirac bi-spinor, and the matrices $\hat{\alpha}_k$ and $\hat{\beta}$ are defined by the following commutational relations:

$$\begin{aligned} \hat{\alpha}_i \hat{\alpha}_k + \hat{\alpha}_k \hat{\alpha}_i &= 2\delta_{ik}, \hat{\alpha}_i \hat{\beta} + \hat{\beta} \hat{\alpha}_i = 0, \\ \hat{\beta}^2 &= 1, i, k = 1, 2, 3 \end{aligned} \quad (2)$$

Let us introduce an evolution operator for the Dirac

equation (1) in a time interval $[t', t'']$:

$$\hat{U}(t'', t') \equiv T \exp \left(-\frac{i}{\hbar} \int_{t'}^{t''} \hat{h} dt \right) \quad (3)$$

Here, T denotes the time ordering of the exponent. The problem is how to represent the kernel of this operator, $\langle x'' | \hat{U} | x' \rangle \equiv \hat{K}(x'', t''; x', t')$, as an integral over trajectories in the 3D configuration space of an electron. Let us notice that this kernel is a 4×4 -matrix in the Dirac bi-spinor space. Below this kernel will be called “propagator”.

Following the standard prescription³, let us begin with the definition of the integral over trajectories in the phase space. Here, we meet the following problem. In the non-relativistic mechanics the Hamiltonian $\hat{h}(p, x)$ is quadratic with respect to the canonical momenta. As a consequence the transition to the path integral in the configuration space is not difficult. Contrary to this case, the Hamiltonian of the Dirac equation is a linear matrix operator with respect to the canonical momenta. A formal integration over the canonical momenta in this case leads to a meaningless expression $\delta^3(\dot{x}^i - c\hat{\alpha}^i)$. In order to overcome this difficulty, we will regularize the path integral in the phase space, adding “by hands” a quadratic form of the canonical momenta. In account of this, let us start with the building of the Feynman path integral.

Let us divide the time interval $[t', t'']$ into N parts $\Delta t_k \equiv [t_k, t_{k-1}]$ of equal length $\varepsilon = (t'' - t')/N$, where $k = 1, \dots, N$, $t_0 \equiv t', t_N \equiv t''$. Basic trajectories in the phase space are broken lines in the 3D-configuration space with pieces $\Delta x_{(k)}^i \equiv x_{(k)}^i - x_{(k-1)}^i$, where we define $x_{(0)}^i \equiv (x^i)'$, $x_{(N)}^i \equiv (x^i)''$. The momentum in any time interval Δt_k has a constant value $p_{i(k)}$.

The standard procedure of iterations leads to the following representation for the propagator of the Dirac

equation (1) in terms of an integral in the phase space:

$$\begin{aligned} \hat{K} = & \lim_{\substack{N \rightarrow \infty \\ \sigma \rightarrow 0}} \int \frac{d^3 p_{(N)}}{(2\pi\hbar)^3} \prod_{k=1}^{N-1} \times \\ & \times \frac{d^3 x_{(k)} d^3 p_{(k)}}{(2\pi\hbar)^3} \times \\ & \times \exp \left[-\frac{\sigma p_{(k)}^2}{2} + \frac{i}{\hbar} p_{i(k)} \Delta x_{(k)}^i \right] \times \\ & \times T \prod_{k=1}^N \left[1 - \frac{i}{\hbar} \varepsilon \hat{h}(p_{(k)}, \tilde{x}_{(k)}) \right] \end{aligned} \quad (4)$$

Here $\tilde{x}_{(k)}^i \equiv (x_{(k)}^i - x_{(k-1)}^i)/2$, and

$$\begin{aligned} \hat{h}(p_{(k)}, \tilde{x}_{(k)}) \equiv & c\hat{\alpha}_i \left(p_{i(k)} - \frac{e}{c} A_i(\tilde{x}_{(k)}) \right) + \\ & + \hat{\beta} m c^2 + e\varphi(\tilde{x}_{(k)}) \end{aligned} \quad (5)$$

The latest multiplier in the Eq.(4) is the time ordered product.

In account of the regularizing quadratic form, the integration over the canonical momenta may be easily performed. As a result, we obtain an integral over trajectories in the configuration space:

$$\begin{aligned} \hat{K} = & \lim_{\substack{N \rightarrow \infty \\ \sigma \rightarrow 0}} \frac{1}{(2\pi\hbar\sigma\varepsilon)^{3/2}} \int \prod_{k=1}^{N-1} \frac{d^3 x_{(k)}}{(2\pi\hbar\sigma\varepsilon)^{3/2}} \times \\ & \times \exp \left(-\frac{\Delta x_{(k)}^2}{2\hbar^2\sigma\varepsilon} \right) \times \\ & \times T \prod_{k=1}^N \left[1 - \frac{i}{\hbar} \varepsilon \hat{h} \left(\frac{i}{\hbar} \frac{\Delta x_{(k)}}{\sigma\varepsilon}, \tilde{x}_{(k)} \right) \right] \end{aligned} \quad (6)$$

This representation of the propagator is convenient for the formulation of the perturbation theory.

First of all, let us demonstrate that in the zero order of the electron charge the integral in Eq.(6) is equal to the well known propagator of a free electron. To this end, let us introduce a momentum representation of the propagator:

$$\begin{aligned} K(\Delta t, p'', p') = & \int d^3 x'' \int d^3 x' \hat{K}(x'', t''; x', t') \times \\ & \times \exp \left(-\frac{i}{\hbar} p'' x'' + \frac{i}{\hbar} p' x' \right) \end{aligned} \quad (7)$$

The integration over coordinates of the end points of a trajectory in Eq.(7) allows to perform the following transformation of the integration variables in Eq.(6): $x_{(k)}^i \rightarrow \Delta x_{(k)}^i, i = 1, \dots, N$, where we take into account that

$$(x^i)'' = (x^i)' + \sum_{k=1}^N \Delta x_{(k)}^i \quad (8)$$

After this, the integral in Eq.(6) may be easily calculated. The integration over $(x^i)'$ gives a multiplier $(2\pi\hbar)^3 \delta^3(p'' - p')$. The integral over $\Delta x_{(k)}^i$ is equal to

$$\begin{aligned} & \prod_{k=1}^N \hbar^3 (2\pi\sigma\varepsilon)^{3/2} \exp \left(-\frac{\sigma\varepsilon (p'')^2}{2} \right) \times \\ & \times \left[1 - \frac{i}{\hbar} \hat{h}(p'') \right] \end{aligned} \quad (9)$$

Here,

$$\hat{h}(p'') \equiv c\hat{\alpha}_k p_{(k)}'' + \hat{\beta} m c^2 \quad (10)$$

is the Hamiltonian of a free electron. The time order of the matrix multiplies in Eq.(9) is unessential. Combining these results, one can obtain the propagator of a free electron:

$$\begin{aligned} \hat{K}^{(0)}(\Delta t; p'', p') = & (2\pi\hbar)^3 \delta^3(p'' - p') \times \\ & \times \exp \left(-\frac{i}{\hbar} \hat{h}(p'') \Delta t \right) \end{aligned} \quad (11)$$

A contribution of the first order in the electric charge to the propagator is equal to

$$\begin{aligned} \hat{K}^{(1)}(t'', t'; p'', p') = & ie \int_{t'}^{t''} dt \exp \left(-\frac{i}{\hbar} \hat{h}(p'') (t'' - t) \right) \times \\ & \times \hat{A}(t, p'' - p') \times \\ & \times \exp \left(-\frac{i}{\hbar} \hat{h}(p') (t - t') \right) \end{aligned} \quad (12)$$

Here

$$\hat{A}(t, q) \equiv \int d^3 x \exp \left(-\frac{i}{\hbar} q x \right) [\hat{\alpha}_k A_k(t, x) - \varphi(t, x)] \quad (13)$$

is the momentum representation of an external electromagnetic field contribution.

In conclusion, using a regularizing quadratic form of the electron momentum in the phase space, we have obtained a representation of the one-electron relativistic quantum dynamics in terms of a path integral over trajectories in the 3D-configuration space. This representation may be used as a basis for the formulation of the relativistic quantum dynamics of many-electron systems with the direct interaction.

* Electronic address: alex.lukyan@rambler.ru

† Electronic address: inna.lukyan@mail.ru

¹ R.P. Feynman. Physics Today, 19, 31, 1966.

² J.D. Bjorken, S.D. Drell, Relativistic Quantum Mechanics,

McGraw-Hill Book Company, N.-Y., 1964.

³ L.D. Faddeev, A.A. Slavnov, Gauge fields, introduction to quantum theory, Benjamin-Cummings, London, 1980.